

# Effect of Purwoceng Extracts (Pimpinella Alpina Molk) on Erectile Function Biomarker Enhancement

*by* Ignatius Riwanto

---

**Submission date:** 31-Jan-2019 12:17PM (UTC+0700)

**Submission ID:** 1070942133

**File name:** Artikel\_C14\_Effect\_of\_Purwoceng\_extract....pdf (253.89K)

**Word count:** 3916

**Character count:** 21242

## Effect of Purwoceng Extracts (*Pimpinella Alpina* Molk) on Erectile Function Biomarker Enhancement

MEINY SUZERY<sup>1</sup>, TAUFIQ UR ACHMAN<sup>2</sup>, IGNATIUS RIWANTO<sup>3</sup>, SUSILO WIBOWO<sup>4</sup>, HADIYANTO<sup>5</sup>

### ABSTRACT

Purwoceng (*Pimpinella alpina* Molk) extract could improve the testosterone levels, LH, and FSH in *Sprague Dawley* male rats. We hypothesized that Purwoceng extract can increase erectile-function biomarker in male rats *Sprague Dawley* through increased expression level of eNOS endothelial, nNOS nerve ending expression levels and the levels of cGMP in penile *corpus cavernosum*. We used Post test only controlled group design and used 30 male rats *Sprague Dawley* which randomized into two groups: castrated and noncastrated, consist of control group, 4 hours and 30-days treatment. Purwoceng extract supplementation dose of 50 mg increased significantly the number of eNOS, nNOS expression and cGMP levels of *corpus cavernosum* penile tissue compare to control group. The highest enhancement of cGMP level along with eNOS and nNOS expression were obtained after 30-days treatment. Enhancement of eNOS expression for endothelium was positively correlated to increment levels of cGMP and nNOS nerve ending smooth muscle *corpus cavernosum*. Purwoceng (*Pimpinella alpina* Molk) extracts could enhance erectile function biomarker.

**Keywords:** Purwoceng, *Pimpinella alpina* Molk, erectile function, cGMP, eNOS, nNOS

### INTRODUCTION

The use of synthetic oral medication for the erectile dysfunction (ED) treatment of sildenafil (Viagra), vardenafil (Levitra) and tadalafil (Cialis) type has been widely used since the '90s. Synthesis of sildenafil derived compounds was done through a series of reactions which quite complicated and selective, starting with diketoster condensation with hydrazine, followed by six other reaction stage<sup>1</sup>. This situation is thought to be the main cause of the high cost of synthetic drugs. In addition, the side effects such as headache, nausea and the decreasing of selectivity to against other PDE isoenzymes, especially PDE-1 and PDE-6, led the researchers interested to develop drugs from natural products. This situation also is supported by the existence of public confidence that natural ingredients are safer for health<sup>2</sup>.

Systematic research of traditional medicine for erectile dysfunction drugs was generally done through the development of aphrodisiac claims<sup>3</sup>. Aphrodisiac is a stimulant substance which can

increase sexual desire by stimulating the excretion of testosterone and accelerate blood circulation to the brain that stimulates nerve function. Some drugs from natural ingredients that had been proven to have an aphrodisiac properties can increase the erection, for example *Tribulus terrestris* and *Panax ginseng*. *Tribulus terrestris* L., contains phytochemicals, flavonoids, alkaloids, glucosides, steroid and saponins. Protodiosin (saponin) was the most dominant compound in this plant, which is a phytochemical agent that can increase sexual arousal and erection through the conversion of testosterone to DHEA, thus can improve erectile dysfunction<sup>4</sup>. In addition, the proerectile has a relaxant effect which is claimed to be an aphrodisiac<sup>5</sup>. *Panax ginseng* root extract contains triterpene saponins called ginsenosides compound which has been reported to improve erectile function by increasing the production of NO. Clinical tests of this plant material also had been reported to have a good effect compared to the placebo<sup>6</sup>. The influence of ginsenosides in raising the nitric oxide (NO) production was done by relaxing the penile smooth muscle through the NO-cGMP pathway<sup>7,8</sup>.

Purwoceng (*Pimpinella alpina* Molk), a typical plant thrives in Indonesia on the Dieng Mountains, Wonosobo, Central Java, have been proven to increase levels of testosterone. This plant has been used for generations to enhance vitality and sexual desire<sup>9</sup>. Purwoceng extract with 50mg dose in male *Sprague Dawley* rats have been proven to increase levels of testosterone, Luteinizing Hormone (LH) and Follicle Stimulating Hormone (FSH)<sup>10</sup>. Phytochemistry

<sup>1</sup>Organic Chemistry Laboratory, Department of Chemistry, Faculty of Science and Mathematic, Diponegoro University, Jl. Prof Soedharto, Tembalang, Semarang

<sup>2</sup>Biochemistry Division and Andrology, Faculty of Medicine, Unisula, Jl. Raya Kaligawe Km 4 Semarang.

<sup>3</sup>Sub Division Digestif Surgery and Doctoral Program, Faculty of Medicine Diponegoro University

<sup>4</sup>Sub Division Biology and Andrology and Post Doctoral Program, Faculty of Medicine Diponegoro University

<sup>5</sup>Chemical Engineering Department, Faculty of Engineering, Diponegoro University,

Correspondence to Meiny Suzery Email: meiny\_suzery@yahoo.com and meinysuzery10@gmail.com

screening of this plant contained steroid, triterpenoid, alkaloid and flavonoid compounds. Analysis of chemical component by gas chromatographic/mass spectroscopy technique of Purwoceng root had showed contained steroid substance as stigmasterol<sup>11</sup>. These compounds from *Pimpinella alpina* Molk were the most playing role in increasing the blood serum's testosterone level and vitality in male rats *Sprague Dawley*.

Study of Purwoceng extract effect on increasing the erectile function biomarkers (eNOS endothelial, nNOS nerve ending and cGMP) has not been studied. Therefore, this study was purposed to prove the effect of Purwoceng extract which can increase the erectile-function biomarker of male rats *Sprague Dawley* through increased expression level of eNOS endothelial, nNOS nerve ending expression levels and the levels of cGMP in penile *corpus cavernosum*.

## MATERIALS AND METHODS

**Materials:** male *Sprague Dawley* rats, materials for making tissue and immunohistochemically examination: formalin solution 10%, Sodium phosphate buffer, xylol, alcohol absolute, alcohol 90% and 70%, paraffin, object glass, antibody monoclonal IgG1 anti-eNOS (1:500) and anti-nNOS (1:500) (oncogene research Products, USA), phosphate buffered solution (PBS), avidin-biotin, biotinylated horseradish peroxidase, mouse monoclonal Ig anti eNOS, anti rat Ig-HRP dan anti-goat Ig-HRP (1:2000), cGMP kit. (Kat#K372-100).

**Plant material and extract:** Root of Purwoceng (*Pimpinella alpina* Molk) were collected in April 2015 from Dieng Mountains, Wonosobo, Central Java, were dried in an oven with circulating air at 40°C. Samples have been deposited in the Herbarium of the "University of Diponegoro" Semarang, Indonesia. Crude extracts were prepared by maceration of 500 g dried Purwoceng roots in room temperature with ethanol 96% (v/v) solvent during 3 x 24 h. The obtained extract was filtered and concentrated in a rotary evaporator at 45°C under reduced pressure (15 mmHg), yielding 5.75 g.

**Animals and Ethics:** *Sprague Dawley* male rat (weight 200-250g) were obtained from Integrated

Research and Testing Laboratory (LPPT) Unit IV, Gadjah Mada University. All the animals were maintained under standard condition, with temperature at 25±2 °C, 12 h dark-light cycle and relative humidity. The rat received food and water ad libitum. This study was approved by the Ethics Committee for Health Research and Medical University School of Medicine and Kariadi Hospital Semarang, Indonesia (Medical Letter No: 016/EC/FK/RSDK).

**Experimental Design:** After acclimatization period, 30 male *Sprague Dawley* rats were randomly allocated into two groups castrated (K) and noncastrated (NK). Each group was divided into a control group, treatment group of 4 hours and 30-days. Purwoceng extract was given orally for the treatment groups by a dose of 50 mg. The treatment was given daily at 8-9 am. The control group only had distilled water. After 4 hours and 30-days all rats were terminated and then the penile tissue was taken for analysis. Expression of eNOS endothelium cell and nNOS nerve ending smooth muscle corpus cavernosum were measured by IHC staining with monoclonal antibody eNOS and nNOS. Measurement of cGMP levels performed by the method of immuno assay on penis tissue supernatant. This procedure was conducted according to Park research<sup>12</sup>.

**Statistical analysis:** The data are expressed as mean ± S.D. Results were statistically evaluated by using one-way analysis of variance (ANOVA) and Turkey's test were carried out to statistically compare the data among the groups. The values were considered significant when p values ≤ 0.05. Regression correlation analysis was used to analyze of data, will be significant if p values ≤ 0.05 with 95% CI.

## RESULTS

The results of assessment of the average amount of eNOS, nNOS and cGMP expression both in castrated and non castrated mice which is performed with period for 4 hours and 30 days shown in Table 1.

Table 1. The average amount of eNOS, nNOS expression and cGMP level in various variables of study

Variables	Treatment group Mean and SD						p
	NK <sub>(-)</sub>	NK <sub>4</sub>	NK <sub>30</sub>	K <sub>(-)</sub>	K <sub>4</sub>	K <sub>30</sub>	
eNOS	6.52±0.60	15.80±0.83	24.80±1.30	5.68±1.17	11.80±0.44	17.84±0.16	0.000 <sup>a</sup>
nNOS	1.08±1.41	4.96±0.16	12.20±0.40	1.84±0.51	4.64±0.29	10.00±0.37	0.000 <sup>a</sup>
Level of cGMP	0.11±0.02	0.15±0.006	3.12±0.38	0.009±0.03	0.13±0.01	0.86±0.34	0.000 <sup>a</sup>

<sup>a</sup>Kruskal Wallis Value p=0.000.



Table 2: Summary of correlation test analysis results

No	Variables	Level of cGMP	
		Correlation Coefficient	p
1	eNOS on endothelium cell CC	0.885	0.000
2	nNOS in smooth muscle CC nerve ending	0.858	0.000

Spearman correlation test  $p < 0.005$

The Mann Whitney test was performed to see whether the difference in eNOS, nNOS expression amount is significant or not on noncastrated rats group compare to controls of treatment for 4 hours and 30 days, the comparison treatment for 4 hours and 30 days showed a high significant difference ( $p < 0.001$ ). Figure 1 and 2 showed the boxplot diagram of eNOS and nNOS expressions.

Fig. 1: Boxplot diagram the number of eNOS expression on endothelium






Fig. 2: Boxplot diagram the number of nNOS nerve ending corpus cavernosum



The statistics tests showed that the purwoceng extract was able to increase the amount of eNOS and nNOS expression in both groups of non-castrated as well as castrated rats. The measurements of the the

mean levels of cGMP in the non-castrated groups showed that administration of the purwoceng extract for 30 days showed highest levels, the difference of each treatment group was analysed by Mann Whitney test, it showed a highly significant difference ( $p < 0.001$ ), whereas the castrated group treated for 4 hours with the controls did not show significant difference ( $p > 0.001$ ).


Fig. 3: Boxplot diagram the level of cGMP on penile tissue



Histopathologic features of eNOS and nNOS expressions with IHC method using a monoclonal antibody staining anti-eNOS and nNOS on mouse tissue for treatment 4 hours and 30 days were shown in Figures 4.

Figure 4 eNOS expression for non-castrated animal group with immunohistochemistry method using antibody monoclonal anti-eNOS coloration (*Mouse monoclonal* IgG1 anti eNOS) on rat penile tissue with 4 hours and 30 days treatment

Fig. 4: eNOS expression for non-castrated animal group with immunohistochemistry method using antibody monoclonal anti-eNOS coloration (*Mouse monoclonal IgG1 anti eNOS*) on rat penile tissue with 4 hours and 30 days treatment



The results showed that the treatment group for 4 hours and 30 days showed the increasing of expression number compared to the control group. The 30-day treatment group showed the highest expression of both treatments. Pathogenesis overview of Purwoceng extract role in improving the erectile function biomarker done through modeling using the line equation, firstly the analysis of correlation tests was performed. The analysis results obtained was a summary as in Table 2.

Based on the correlation test results show that the two variables play role in increasing the levels of cGMP.

Multivariate analysis with Spearman test obtained the regression line of the effects of purwoceng extract on the levels of cGMP are follows:  $0.923 + 0.109$  (nNOS nerve endings) +  $0.76$  (eNOS endothelium). This equation indicates that the Purwoceng extract treatment contribute to the changes in the levels of cGMP along with the increasing number of eNOS and nNOS expression.

## DISCUSSION

Research variables are the expression of eNOS, nNOS and cGMP levels in penile tissue of rats. The

three dependant variable is directly related to NO, because NO is produced through a reaction catalyzed by specific enzyme NOS<sup>13</sup>. Nitric Oxide is released from nerve endings nitregik produced by oxidation of L-arginine catalyzed by nNOS, whereas NO released from endothelial cells is catalyzed by eNOS. As a major neurotransmitter, NO is responsible for the occurrence of smooth muscle relaxation in the corpus cavernosum and diffuses directly into the corpus cavernosum smooth muscle guanylyl cyclase to activate enzymes to accelerate the formation of cGMP that causes relaxation of corpus cavernosum smooth muscle. The cGMP equilibrium is regulated by PDE-5 enzyme to change cGMP into 5-GMP. Thus increased levels of eNOS and nNOS would lead to an increase in cGMP levels, so these three variables is very important in the process of relaxation in the corpus cavernosum smooth muscle of the penis. Therefore, eNOS, nNOS and cGMP are biomarkers of erectile function.

Endothelial NOS is an enzyme the identified attached to the membrane of blood vessels and serve as material for vasodilation. NO produced by eNOS is responsible for keeping blood vessels functioning normally and as a mediator of the erectile mechanism<sup>14,15,16</sup>. The results showed increament of the number of eNOS expression in Purwoceng extract for 4 hours and 30 days were significant compared to the control group rats castrated and non-castrated. Thus, increasing the amount of eNOS expression is also assumed to increase NO production. Increasing the amount of expression is highly influenced by the hormone testosterone, because testosterone is a central mechanism to regulate the erection process. In this study, enhancement of the amount eNOS expression allegedly caused by the presence of elevated levels of serum testosterone. Evidence showed that the effect of extract Purwoceng increase serum levels of testosterone through LH simulation and through the conversion of stigmasterol to testosterone in peripheral tissues<sup>10</sup>. The presence of stigmasterol compound in the extract Purwoceng also been shown<sup>11</sup>.

This study uses the castrated rats, are intended to describe the state of andropouse because castrated can cause loss of testosterone-producing Leydig cells. While the use of non-castrated rat meant to portray a normal state. The results showed that the effect of administration Purwoceng extract in the castrated groups of rats for 4 hours and 30 days compared to normal rat groups showed differences in the amount of eNOS expression is significantly ( $p < 0.001$ ). This is due to an increase in the amount of eNOS expression in castrated rats, because there is



the adrenal cortex to produce testosterone, although in very small amounts<sup>17,18,19,20</sup>.

Provision for 30 days showed the highest expression of eNOS compared to 4 hours of treatment and control groups. This suggests that the longer time of administration, the higher expression of eNOS. These results reinforce earlier theory that medicinal plants takes time to reach and influence the target organ.

Increasing the amount of eNOS expression is suspected by the presence of bioactive flavonoid compounds in extracts Purwoceng. Flavonoid shown induce eNOS, increasing cytosolic calcium that works as a cofactor of eNOS activate and eliminate free radicals in the interstitial fluid that NO production can be maintained. In addition, flavonoids work as vasorelaxant by inhibiting PDE-5 and lower  $Ca^{+2}$  in smooth muscle cells<sup>21</sup>.

Flavonoid compounds have been found in the plant genus *Pimpinella* is illunginin A f B<sup>22</sup>. Effects of flavonoids have also been shown to activate eNOS in the provision of epigallocatechin gallate in the reperfusion of ischemic hearts. Another researcher, reported that injection of resveratrol which is a flavonoid compound intraperitoneally in rat showed increased expression of eNOS and nNOS<sup>23</sup>.

Neuronal NOS is an enzyme located in the nerve tissue and striated muscle, which serves as a communication cell. nNOS in the penis expressed as neurons innervating the corpus cavernosum penis. Neuronal NOS to synthesize NO are dominant in the brain and released from the nerve terminal. Mechanism of action of nNOS in the relaxation process is not yet fully understood. Allegedly NO derived from L-arginine stimulates the activity of guanylyl cyclase to convert GTP to cGMP in the corpus cavernosum smooth muscle of the penis<sup>24,25</sup>.

The results showed that there is a significant increase in the number of nNOS expression in the non-castrated and castrated rats after administration of Purwoceng extract for 4 hours and 30 days compared to the negative control group ( $p < 0.001$ ). The number of nNOS expression is proven to increase due to administration of a dose of 50 mg extract Purwoceng. Increasing the number of nNOS expression thus also increases the ability of an erection, because nNOS is a biomarker that erectile function can produce NO. An increase of the number of eNOS and nNOS expression, is expected to increase levels of NO resulting in vasodilation and smooth muscle relaxation in the corpus cavernosum penis of male *Sprague Dawley* rats.

Nitric oxide (NO) the activates the enzyme quanylate cyclase and induces smooth muscle relaxation by increasing cGMP levels. Cyclic GMP will stimulate protein kinase G (PKG) which is

inactive to active. Then the active PKG lowers calcium levels, causing dilatation and relaxation of corpus cavernosum smooth muscle, which in turn causes an erection. Cyclic GMP are second messengers resulting from changes in the GTP activated by NO via activation of guanylyl cyclase enzyme<sup>26</sup>. NO neurotransmitters and second messengers cGMP are both relaxing vascular smooth muscle. Purwoceng extract in castrated and non-castrated for 30 days in this study increase cGMP levels significantly compared to the control group ( $p < 0.001$ ). These results prove that an increase in cGMP levels due to administration of the extract Purwoceng. The increasing of cGMP levels through increased expression of eNOS and nNOS.

Based on the Spearman correlation test of the biomarker increased erectile function it is obtained that the increasing role Purwoceng extract cGMP accompanied by increased expression levels of nNOS and eNOS. The increase of all biomarker eNOS, nNOS and cGMP caused by the presence of elevated levels of serum testosterone.

## CONCLUSION

Purwoceng extract supplementation dose of 50 mg on male *Sprague Dawley* rats among castrated and non-castrated has been proven increased the biomarker erectile function. It has been proven from increase of the number of eNOS expression in endothelial cells, nNOS in the nerve endings smooth muscle of corpus cavernosum penile tissue, and cGMP levels. The highest enhancement were obtained after 30-days treatment. Enhancement of cGMP level was positively correlated to increament eNOS endothelium and nNOS nerve ending smooth muscle corpus cavernosum.

**Conflict of interest:** We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

## REFERENCE

1. Terret, N.K., Bell, A.K., Brown, D., & Ellis, P., 1996. *Bioorg. Med. Chem. Lett.*, 6, pp.1819–1824.
2. Gratz, S.R., Flurer, C.L. & Wolnik, K.A., 2004. N. *J.Pharm. Biomed. Anal.*, 36, pp.525–533.
3. Traish, A.M., Goldstein, I. & Kim, N.N., 2007. Testosterone and Erectile Function: From Basic Research to a New Clinical Paradigm for Managing Men with Androgen Insufficiency and Erectile Dysfunction. *European Urology*, 52(1), pp.54–70.
4. Adimoelja, A., 2000. Phytochemicals and the breakthrough of traditional herbs in the management of sexual dysfunctions. *International Journal of Andrology*, 23(S2), pp.82–84.
5. Andaikan, P.G., Gauthaman, K. & Prasad, R.N., 2000. Pro-erectile Pharmacological Effects of Tribulus Terrestris Extract on The Rabbit Corpus Cavernosum. *Ann. Acad. Med. Singapore*, 29,

- pp.22–26.
6. Choi, H., Seong, D. & Rha, K., 1995. Clinical Efficacy of Korean Red Ginseng for Erectile Dysfunction. *Int J Impot Res*, 7, pp.181–186.
7. Kim, Ji H., Hahm, D. H., Yang, D. C., Kim, Jang H., Lee, H. J., & Shim, Insop, 2005. Effect of Crude Saponin of Korean Red Ginseng on High-Fat Diet- Induced Obesity in the Rat. *Journal of Pharmacological Sciences*, 97, pp.124–131.
8. Young-Mi, K., Seung, N., Yun, Y. G., Hong, H. D., Lee, Y. C., Ha, K. S., Lee, H., Kwon, H. J., Kwon, Y. G., & Kim, Y. M., 2007. Water Extract of Korean Red Ginseng Stimulates Angiogenesis by Activating the PI3K/Akt-Dependent ERK1/2 and eNOS Pathways in Human Umbilical Vein Endothelial Cells. *Biol. Pharm. Bull*, 30(9), pp.1674–1679.
9. Caroeboka, A.M. & Lubis, I., 1975. Pemeriksaan Pendahuluan Kandungan Kimia akar Pimpinella alpina dalam Simposium Tanaman Obat I., pp.Bogor, 8–9 Desember.
10. Taufiqurrachman & Wibowo, S., 1999. Pengaruh Ekstrak Pimpinella alpina Molk dan Eurycoma Longifolia Jack terhadap Peningkatan Kadar Testosteron LH dan FSH pada Tikus Jantan Spraque Dawley. *Media Medika Indonesia*, 35(2).
11. Suzery, M., Ngadiwiyono, Nurhasnawati, E., & Cahyono, B., 2004. Senyawa Stigmasterol dari Pimpinella alpina Molk. *Media Medika Indosiana*, 29 Suppl.(1), pp.37–39.
12. Park, C.S., Ryu, S.D. & Hwang, S.Y., 2004. Herbal Formula Enhances Erectile Responses in Aged, Diabetic, and Normal Rats via the Activation of NO/cGMP Signaling Pathways. *Journal of Ethnopharmacology*, 94, pp.85–92.
13. Moncada, S. & A, H., 1993. The L-arginine-nitric Oxide Pathway. *N. Engl. J. Med.*, 329(27), pp.2002–2012.
14. Moncada, S. & Higgs, A., 1997. The Arginine-nitric Oxide Pathway. *N. Engl. J. Med.*, 329, pp.2002–2012.
15. Ghalayini, I., 2004. Nitric Oxide-cyclic GMP Pathway with Some Emphasis on Cavernosal Contractility. *International Journal of Impotence Research*, 16, pp.459–469.
16. Moncada, S. & Higgs, A., 2006. The Discovery of Nitric Oxide and Its Role in Vascular Biology. *Br. J. Pharmacol*, 147(Suppl. I), pp.S193–201.
17. Cooper, T.G. & Yeung, C.H., 1977. Physiology of Sperm Maturation and Fertilization in: Neischlag E, Behre AM, editors *Andrology, Male Reproduction Health and Dysfunction. Berlin Heldeberg: Springer-Verlag*, pp.178–181.
18. Van, A.H. & Hertle, R., 1997. Erectile Dysfunction, Disorder of Sperm Deposition in: Neischlag E, Behre HM, (Editions), *Andrology, Male Reproductive Health and Dysfunction. Berlin Heldeberg: Springer-Verlag*, pp.178–181.
19. Howel, S. & Shalet, S., 2001. Testosterone Deficiency and Replacement. *Horm. Res.*, 56(1), pp.82–92.
20. Thijssen, J.H.H., 2002. Laboratory Test in Endocrine Evaluation of Aging Male (Appendix) in: Lunenfeld B, Gooren L (editors), *Textbook of Men's Health*, London: Yhe Pathenon Publishing Groups., pp.44–50.
21. Akhlaghi, M. & B., B., 2009. Mechanisms of Flavonoid Protection Against Myocardial Ischemia-reperfusion Injury. *Journal of Molecular and Cellular Cardiology*, 46, pp.309–317.
22. Qiao, B.-L., Wang, C. D., Shi, H., Mi, C. F., & Li, F., 1996. Studies on Chemicals Constituent of the Root of Pimpinella Thellungiana. *Zhongcaoyao*, 27(3), pp.136–138.
23. Hung, L.M., Su, M.J. & Chen, J.K., 2004. Resveratrol Protects Myocardial Ischemia-reperfusion Injury Throught Both NO-dependant and NO-independant Mechanisms. *Free Radic. Biol. Med.*, 36, pp.774–781.
24. Magee, T.R., Ferrini, M. G., Davila, H. H., Zeller, C. B., Vernet, D., Sun, J., Lalani, R., Burnett, A. L., Rajfer, J., & Gonza, N. F., 2003. Protein Inhibitor of Nitric Oxide Synthase (NOS) and the N-Methyl-D-Aspartate Receptor Are Expressed in the Rat and Mouse Penile Nerves and Colocalize with Penile Neuronal NOS. *BIOLOGY OF REPRODUCTION*, 68, pp.478–488.
25. Toda, N., Ayajiki, K. & Okamura, T., 2005. Nitric oxide and penile erectile function. *Pharmacology and Therapeutics*, 106(2), pp.233–266.
26. ucas, K., Pitari, G.M., Kazerounian, S., Ruiz-Stewart, I., Park, J., & Schulz, S., 2000. Guanylyl Cyclases and Signaling by Cyclic GMP. *Pharmacol Rev.*, 52, pp.375–414.

# Effect of Purwoceng Extracts (Pimpinella Alpina Molk) on Erectile Function Biomarker Enhancement

## ORIGINALITY REPORT

24%

SIMILARITY INDEX

18%

INTERNET SOURCES

20%

PUBLICATIONS

14%

STUDENT PAPERS

## PRIMARY SOURCES

1

Submitted to University of Warwick

Student Paper

2%

2

M Suzery, B Cahyono, P Astuti. " The Acute Toxicity Test of Methanolic Extract of Poit on Liver Balb/c Mice ", IOP Conference Series: Materials Science and Engineering, 2017

Publication

2%

3

[onlinelibrary.wiley.com](http://onlinelibrary.wiley.com)

Internet Source

1%

4

[theses.gla.ac.uk](http://theses.gla.ac.uk)

Internet Source

1%

5

Submitted to University College London

Student Paper

1%

6

Submitted to University of New England

Student Paper

1%

7

Li, X.. "Anticipatory postural adjustments in conditions of simulated reduced gravity", Gait & Posture, 200811

1%



8

C.H. Yeung. "Human sperm volume regulation. Response to physiological changes in osmolality, channel blockers and potential sperm osmolytes", Human Reproduction, 05/01/2003

Publication

1 %

9

Ana Pavla A. Diniz Gurgel, Jackeline G. da Silva, Ana Ruth S. Grangeiro, Danielli C. Oliveira et al. "In vivo study of the anti-inflammatory and antitumor activities of leaves from *Plectranthus amboinicus* (Lour.) Spreng (Lamiaceae)", Journal of Ethnopharmacology, 2009

Publication

1 %

10

[www.uaiasi.ro](http://www.uaiasi.ro)

Internet Source

1 %

11

Submitted to South Bank University

Student Paper

1 %

12

[tees.openrepository.com](http://tees.openrepository.com)

Internet Source

1 %

13

Sohn, D.W.. "Elevation of intracavernous pressure and NO-cGMP activity by a new herbal formula in penile tissues of spontaneous hypertensive male rats", Journal of Ethnopharmacology, 20081120

Publication

1 %

14	Pacanoski, Zvonko, Štefan Týr, and Tomáš Vereš. "Puncturevine (Tribulus terrestris L.): noxious weed or powerful medical herb", Journal of Central European Agriculture, 2014. Publication	1 %
15	Submitted to University of Newcastle Student Paper	1 %
16	Submitted to University of Leicester Student Paper	1 %
17	Submitted to University of Reading Student Paper	1 %
18	Submitted to Imperial College of Science, Technology and Medicine Student Paper	1 %
19	www.auanet.org Internet Source	1 %
20	Gurgel, A.P.A.D.. "In vivo study of the anti-inflammatory and antitumor activities of leaves from Plectranthus amboinicus (Lour.) Spreng (Lamiaceae)", Journal of Ethnopharmacology, 20090907 Publication	<1 %
21	ejournal.undip.ac.id Internet Source	<1 %

22

Internet Source

&lt;1 %

23

[edocs.fu-berlin.de](http://edocs.fu-berlin.de)

Internet Source

&lt;1 %

24

Walter Krause. "Testosterone supplementation in the aging male: Which questions have been answered?", *The Aging Male*, 3/2005

Publication

&lt;1 %

25

[www.jstage.jst.go.jp](http://www.jstage.jst.go.jp)

Internet Source

&lt;1 %

26

[www.jbc.org](http://www.jbc.org)

Internet Source

&lt;1 %

27

Marques, Davi Lazarini, Mônica Alves, Carolina Maria Modulo, Lilian Eslaine Costa Mendes da Silva, Peter Reinach, and Eduardo Melani Rocha. "Lacrima osmolarity and ocular surface in experimental model of dry eye caused by toxicity", *Revista Brasileira de Oftalmologia*, 2015.

Publication

&lt;1 %

28

[iovs.arvojournals.org](http://iovs.arvojournals.org)

Internet Source

&lt;1 %

29

Maria Galati, Enza, and Jurgen Reichling. "Chemical Constituents of the Genus *Pimpinella*", *Illicium Pimpinella* and

&lt;1 %



30

Paola Ascoli. "Hypopituitarism", Pituitary, 12/2006

Publication

<1 %

31

[www.termedia.pl](http://www.termedia.pl)

Internet Source

<1 %

32

[aaps.org](http://aaps.org)

Internet Source

<1 %

33

[www.bioone.org](http://www.bioone.org)

Internet Source

<1 %

34

El Maghraby, Gamal M., Fars Kaed Alanazi, and Ibrahim A. Alsarra. "Transdermal Delivery of Tadalafil. I. Effect of Vehicles on Skin Permeation", Drug Development and Industrial Pharmacy, 2009.

Publication

<1 %

35

"Sexual Function in the Prostate Cancer Patient", Springer Nature America, Inc, 2009

Publication

<1 %

36

Kazuhiko Komori. "Nitric Oxide Synthesis Leads to Vascular Endothelial Growth Factor Synthesis via the NO/Cyclic Guanosine 3',5'-Monophosphate (cGMP) Pathway in Human Corpus Cavernosal Smooth Muscle Cells", The Journal of Sexual Medicine, 3/4/2008

<1 %

---

37	<a href="http://neuronresearch.net">neuronresearch.net</a>	<1 %
	Internet Source	

---

38	<a href="http://www.nature.com">www.nature.com</a>	<1 %
	Internet Source	

---

39	Muller, M.. "Sex hormones and male health: effects on components of the frailty syndrome", Trends in Endocrinology & Metabolism, 200308	<1 %
	Publication	

---

40	<a href="http://citeseerx.ist.psu.edu">citeseerx.ist.psu.edu</a>	<1 %
	Internet Source	

---

41	Enrico de Andrade. "Study of the efficacy of Korean Red Ginseng in the treatment of erectile dysfunction", Asian Journal of Andrology, 3/2007	<1 %
	Publication	

---

42	Yuan, Xiao, Xueling Guo, Yan Deng, Die Zhu, Jin Shang, and Huiguo Liu. "Chronic intermittent hypoxia-induced neuronal apoptosis in the hippocampus is attenuated by telmisartan through suppression of iNOS/NO and inhibition of lipid peroxidation and inflammatory responses", Brain Research, 2015.	<1 %
	Publication	

---

43	<a href="http://mscanner.stanford.edu">mscanner.stanford.edu</a>	
----	--	--

&lt;1 %

44

[www.yourbrainonporn.com](http://www.yourbrainonporn.com)

Internet Source

&lt;1 %

45

"Contemporary Treatment of Erectile Dysfunction", Springer Nature America, Inc, 2011

Publication

&lt;1 %

46

[medlibes.com](http://medlibes.com)

Internet Source

&lt;1 %

47

Jung, Ae Ryang, Yong Sun Choi, Shuyu Piao, Yong Hyun Park, Kshitiz Raj Shrestha, Seung Hwan Jeon, Sung-Hoo Hong, Sae Woong Kim, Tae-Kon Hwang, Ki Hean Kim, and Ji Youl Lee. "The Effect of PnTx2-6 Protein From Phoneutria nigriventer Spider Toxin on Improvement of Erectile Dysfunction in a Rat Model of Cavernous Nerve Injury", Urology, 2014.

Publication

&lt;1 %

Exclude quotes

On

Exclude matches

&lt; 1 words

Exclude bibliography

On



# Effect of Purwoceng Extracts (Pimpinella Alpina Molk) on Erectile Function Biomarker Enhancement

## GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

**Instructor**

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6